

PlaqueOff



PRODEN PLAQUEOFF® POWDER

Influence of ProDen PlaqueOff® Powder on




Conclusion

In both conducted studies, ProDen PlaqueOff® Powder containing A.N ProDen® efficiently and significantly decreased plaque and calculus accumulation as well as gingival bleeding. Based on the results of these studies product ProDen PlaqueOff® Powder was awarded by VOHC approval.



Figure 1.

Influence of ProDen PlaqueOff® on plaque, calculus and gingival bleeding, results from study 2.



Objective

To determine the influence of the product ProDen PlaqueOff Powder containing the brown algae A.N ProDen® on plaque, dental calculus accumulation and gingival bleeding index in dogs.

Material and Methods

The studies were carried out in two groups of 30 dogs, male and female, of Cavalier King Charles Spaniel, Miniature Schnauzer, Japanese Spaniel, Corgi, Chihuahua, Yorkshire and mixed breeds. The animals were randomly subdivided into two groups receiving ProDen PlaqueOff® Powder, with A.N ProDen® algae, or placebo (microcrystalline cellulose granules, without the algae).

After 30 days, a clinical evaluation of the plaque index, calculus index, and gingival bleeding index was carried out, taking into account the weight, general appearance, and blood test results of the patients.

Results

After 30 days of oral administration, ProDen PlaqueOff® Powder significantly reduced the accumulation of plaque and calculus as well as reduced the gingival bleeding index. The mean reduction was 32% (plaque index), 35% (calculus index) and 67% (gingival bleeding index).

Table 1. Results on dogs in study 1 0,3852 +/- 0,2417 P1 Group 2,207 +/- 0,5667 0,8778 +/- 0,2414 P2 Group 1,489 +/- 0,5858 0,6259 +/- 0,3863 0,1556 +/- 0,1192 Statistical test Parametric non-paired t-test Parametric non-paired t-test Parametric non-paired t-test YES: 0,0020 YES: 0,0411 YES: 0,0026 Statistical significance (P value) Difference between P2 vs P1 (%) 28% 32% 59%

Table 2. Results on dogs in study 2

	Plaque Index (mean +/- SD)	Calculus score (mean +/- SD)	Gingival bleeding index (mean +/- SD)
P1 Group	2,396 +/- 0,9595	0,9815 +/- 0,5385	0,4111 +/- 0,2847
P2 Group	1,633 +/- 0,4999	0,563 +/- 0,3099	0,1037 +/- 0,1292
Statistical test	Parametric non-paired t-test	Parametric non-paired t-test	Parametric non-paired t-test
Statistical significance (P value)	YES: 0,0108	YES: 0,0144	YES: 0,0007
Difference between P2 vs P1 (%)	31%	42%	74%

Gawor Jerzy, Klinika Arka Krakow, Poland - Michal Jank, Division of Pharmacology and Toxicology; Faculty of Veterinary Medicine, Warsaw University of Life Science, Poland - Katarzyna Jodkowska, Faculty of Veterinary Medicine, Warsaw University of Life Science, Poland - Emilia Klim, Klinika Pulawska Warszawa, Poland - Ulla K Svensson, UKS Life Science Consulting AB, Lund, Sweden